



# Published Opposites Attract Study Supports Preventive Power of Being Undetectable

The study of mixed-HIV-status gay couples prompted the CDC to conclude that undetectable people have effectively no risk of passing on HIV.

July 20, 2018 By [Benjamin Ryan](#)

---

The Lancet has published the study that, paired with a previous, similar study, led the Centers for Disease Control and Prevention (CDC) to conclude in 2017 that people with HIV who have an undetectable viral load have effectively no risk of transmitting the virus. Called Opposites Attract, this observational cohort study of mixed-HIV-status gay male couples was first [presented](#) to great fanfare at the 9th International AIDS Society Conference on HIV Science in Paris (IAS 2017) in July 2017.

In Opposites Attract, the participants contracted sexually transmitted infections (STIs) at a high rate and yet still did not transmit HIV among them. This fact helps challenge the presumption that contracting STIs may raise the risk of HIV transmission even when the partner with the virus has a fully suppressed viral load according to his most recent test.

The study authors still recommend that mixed-HIV-status gay couples use condoms or Truvada (tenofovir disoproxil fumarate/emtricitabine) as pre-exposure prophylaxis (PrEP) during the first six months that the partner with HIV is on treatment for the virus, until tests make certain that he has a fully suppressed viral load.

Opposites Attract included 343 male-male couples in which one partner was HIV positive and the other was HIV negative. The men were recruited at 13 clinics in Australia (from May 2012 to March 2016), one in Brazil (from May 2014 to March 2016) and one in Thailand (also from May 2014 to March 2016). All the couples made at least one follow-up visit and were followed for a cumulative 588.4 years. (This follow-up figure looked at years of follow-up per couple, not per individual participant.)

At their study visits, the HIV-negative partners provided information about their sexual behavior and received HIV and STI testing. The HIV-positive partners received viral load testing (with viral suppression defined as a viral load below 200) as well as CD4 count and STI testing.

If HIV-negative partners contracted the virus, the researchers conducted genetic testing to determine whether their virus matched their partner's virus, thus suggesting that the virus transmitted between them.

The researchers focused their final analysis on periods during which the couples reported condomless anal sex between them, the HIV-negative partner was not using PrEP and the HIV-positive partner was virally suppressed.

Both the HIV-positive and HIV-negative men had an average age of 34. At their initial study visit, 13 percent of the HIV-positive partners had an STI and 80 percent were taking antiretroviral (ARV) treatment. Of the HIV-treated group, 88 percent reported taking more than 90 percent of their ARV doses during the previous three months; 78 percent had a fully suppressed viral load. Of the HIV-negative partners, 8 percent reported recent PrEP use and 11 percent had an STI.

During follow-up, of the HIV-positive partners, 2 percent never took ARVs, 25 percent started ARVs during the study and 73 percent were always on treatment for the virus. Seventy-five percent of the HIV-positive partners consistently had a viral load below 200, 23 percent had a viral load that varied between below and above 200, and 2 percent had a viral load consistently above 200. Thirty-four percent of the HIV-positive partners were diagnosed with an STI. Of the HIV-negative partners, 34 percent reported PrEP use at any time during follow-up; 25 percent were diagnosed with an STI.

A total of 253 (74 percent) of the couples reported condomless anal sex within each couple, for a total of 16,800 condomless sex acts. Three of the HIV-negative partners in these couples contracted the virus during follow-up. None of these viral strains were genetically matched to the respective men's primary partner. Consequently, the researchers concluded that there were zero transmissions within these couples.

There were 232.2 cumulative years of follow-up and 12,447 condomless anal sex acts during periods when partners reported condomless anal sex, the HIV-positive partners were virally suppressed and the HIV-negative partners were not on PrEP. Based on this amount of data, the researchers' statistical analysis led them to conclude that the true risk of HIV transmission in this context was between zero and 1.59 transmissions per 100 cumulative years.

This estimate range is known as a confidence interval. If the study had had more follow-up time and still no HIV transmissions, this confidence interval would have been narrower, with a smaller figure for the upper estimate of the true transmission risk and the low-end of the estimate still zero.

In 2016, the [PARTNER](#) study also reported no linked HIV infections between mixed-HIV-status gay couples, who were followed for a cumulative 415 years during which they reported condomless sex among them, the HIV-positive men were virally suppressed and the HIV-negative men were not on PrEP. The authors of that study estimated that the true risk of infection was between zero and 0.89 infections per 100 years of sex between such couples.

The Opposites Attract study authors wrote, “This study provides evidence that HIV transmission in the context of viral suppression is very low, although we cannot exclude higher levels of risk based on the upper [confidence interval] limit.” They stressed that between PARTNER and Opposites Attract, there have been no linked transmissions of HIV despite nearly 35,000 acts of condomless sex between the gay couples.

Andrew Grulich, head of the Kirby Institute at UNSW Sydney’s HIV Epidemiology and Prevention Program and chief investigator on the study, was much more emphatic in a press release, stating, “These results form a significant part of the evidence base for the international community-led Undetectable=Untransmissible, or U=U campaign, which highlights the fact that people living with HIV can now live long and healthy lives, with effectively zero chance of sexually transmitting the virus to others, provided their viral load is undetectable due to effective ART.”

Andrew Grulich, PhD, an HIV epidemiologist at the Kirby Institute at UNSW Sydney in Australia at IAS 2017 Benjamin Ryan

While the 2011 HPTN 052 study of mixed-HIV-status heterosexual couples saw four genetically matched transmissions within the couples during the first six months that the respective HIV-positive partners were on ARVs, Opposites Attract did not observe this phenomenon. Given the relatively small data set on hand in the latter study when it came to calculating the risk of transmission during this first six-month period of HIV treatment, the study authors stated that “the most appropriate policy and community education response might be to suggest use of condoms

or PrEP in the first six months [of ARV treatment for the HIV-positive partner in a mixed-HV-status couple] and until viral suppression is certain. After this period, viral suppression alone is likely to be sufficient [to prevent HIV transmission].

“However,” the study authors continued, “HIV-negative men in nonmonogamous serodiscordant couples should be considered for PrEP initiation, given the high HIV incidence among these men and that [condomless anal sex] with outside partners was not uncommon.”

The researchers calculated that when the partners used one or more HIV risk reduction strategy—condoms, PrEP or viral suppression—the true risk of transmission between them was between zero and 0.63 transmissions per 100 cumulative years of follow-up. That calculation was based on the 582.5 cumulative years of follow-up during which one or more of these strategies were in place among the couples.

To read a press release about the study, [click here](#).

To read the study abstract, [click here](#).

---

© 2024 Smart + Strong All Rights Reserved.

<https://www.poz.com/article/opposites-attract-draft>