



Viramune and Viramune XR

Generic Name: nevirapine

Pronunciation: VIH-rah-mune

Abbreviation: NVP

Drug Class: [Non-Nucleoside Reverse Transcriptase Inhibitors \(NNRTIs\)](#)

Company: Boehringer-Ingelheim

Approval Status: Approved

Generic Version Available: Yes

Drug Indication

Not part of a recommended or alternative treatment regimen for antiretroviral-naive people living with HIV, according to the U.S. Department of Health and Human Services (DHHS) Panel on Antiretroviral Guidelines for Adults and Adolescents. Visit

http://aidsinfo.nih.gov/contentfiles/lvguidelines/aa_recommendations.pdf for the full DHHS guidelines.

General Info

Viramune is an HIV medication. It is in a category of HIV medicines called non-nucleoside reverse transcriptase inhibitors.

Viramune was approved by the U.S. Food and Drug Administration for use by people living with HIV in 1996.

Viramune is available as an immediate-release (IR) tablets, which is taken twice a day, and as an extended-release tablet (XR), which is taken once a day.

Viramune must be used in combination with other HIV drugs.

Dosage

Adult Dose: One 200mg Viramune IR tablet once a day for 14 days, then one 400mg Viramune XR tablet once a day

Pediatric Dose: Age 6 to 18 years: One 150mg Viramune IR tablet or liquid for the first 14 days. Dosing then based on body surface area.

Dosing Info: Take with or without food.

Side Effects

The U.S. Food and Drug Administration (FDA) recommends that women with CD4 counts over 250 and men with CD4 counts over 400 should not take Viramune as their first antiretroviral (ARV) regimen unless the benefits outweigh the risks. This is because Viramune can lead to serious liver problems in people with higher CD4 counts who start Viramune for the first time. This is especially true in pregnant women with higher CD4 counts. People whose CD4 counts have significantly increased from taking another ARV regimen (to greater than 250 in women and 400 in men) may safely switch to Viramune, however.

In rare cases, liver problems can be very serious and become life-threatening. Therefore, if you develop any of the following symptoms while taking Viramune, call your doctor immediately: general ill feeling, tiredness, lack of appetite, yellowing of the skin or eyes, darkening of the urine, pale stools, liver tenderness.

The most common side effect of Viramune is skin rash. In a small number of patients, rash has been serious and resulted in death. Therefore, if you develop a rash with any of the following symptoms, call your doctor immediately: general ill feeling, fever, muscle or joint aches, blisters, mouth sores, conjunctivitis (inflammation of the inner surface of the eyelids), swelling of the face, tiredness.

If you must stop treatment with Viramune because you have these types of serious reactions, you must not take Viramune again.

Drug Interactions

For a review of drug interactions, including prescription and over-the-counter medications and supplements that should not be taken with Viramune or may require dose adjustments, consult the Viramune package insert:

<http://bidocs.boehringer-ingenelheim.com/BIWebAccess/ViewServlet.ser?docBase=renetnt&folderPath=/Prescribing+Information/PIs/Viramune+XR/ViramuneXR.pdf>

Other Info

Before taking this medication, tell your doctor if you have kidney disease or liver disease. In addition, tell your doctor if you are pregnant or planning to become pregnant, if you are breastfeeding, and all your medical conditions, including all prescription and over-the-counter medications and supplements you are taking.

For More Info:

<http://bidocs.boehringer-ingelheim.com/BIWebAccess/ViewServlet.ser?docBase=renetnt&folderPath=/Prescribing+Information/PIs/Viramune+XR/ViramuneXR.pdf>

Co-Pay Program Info: http://www.aidsmeds.com/articles/PAPs_Copays_19740.shtml

Patient Assistance Program Info: http://www.aidsmeds.com/articles/PAPs_Copays_19740.shtml

Last Reviewed: July 15, 2016

© 2024 Smart + Strong All Rights Reserved.

<https://www.poz.com/drug/viramune-viramune-xr>